

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# Chipco Rovral<sup>®</sup> GT

FUNGICIDE

Active Constituent: 250 g/L IPRODIONE  
 Solvent: 331 g/L LIQUID HYDROCARBONS

GROUP **B** FUNGICIDE

**For control of fungal diseases in ornamentals and recreational turf as specified in the  
 DIRECTIONS FOR USE table**

## GENERAL INSTRUCTIONS

### Fungicide Resistance Strategy

For fungicide resistance management Chipco Rovral GT Fungicide is a Group **B** fungicide. Some naturally occurring individual fungi resistant to Chipco Rovral GT Fungicide and other Group **B** fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Chipco Rovral GT Fungicide and other Group **B** fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience accepts no liability for any losses that may result from the failure of Chipco Rovral GT Fungicide to control resistant fungi.

**Note:** Chipco Rovral GT may be very unstable in conditions where the pH is 7 or higher. It is therefore essential to check the pH of the spray solution before adding Chipco Rovral GT. A suitable registered buffering agent may have to be added to bring pH down below 7.

### Application

Good control requires good coverage. Application should be made using sufficient water to ensure thorough coverage.

### Turf Nutrition

This product acts very rapidly and where the turf is of a low nutritional status, a slight yellowing may be seen. This will be corrected by nitrogenous fertiliser application. Where possible, it is recommended that the fertiliser program should precede the fungicide application.

### Compatibility

With any mixture, thoroughly agitate immediately before applying. The use of a surfactant or spray oil is not recommended with Chipco Rovral GT as it may result in damage to sensitive plants. DO NOT mix with fertilisers. Mixtures with some fertilisers, e.g. urea, may cause foliar damage. Mixing Chipco Rovral GT with Chipco Aliette<sup>®</sup> Signature may result in some settling out.

As formulations of other manufacturers' products are beyond the control of Bayer CropScience, all mixtures should be tested prior to mixing commercial quantities.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on-site. Break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (ph: 131126).

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet.

**EXCLUSION OF LIABILITY**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

Chipco® and Rovral® are Registered Trademarks of Bayer.

NRA Approval No.: 50954/0103

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

**DIRECTIONS FOR USE**

**PRECAUTIONS: DO NOT GRAZE TREATED AREAS OR FEED TURF CLIPPINGS FROM TREATED AREAS TO ANIMALS INCLUDING POULTRY**

SITUATION	DISEASE	RATE	CRITICAL COMMENTS
Ornamentals	Botrytis Blight ( <i>Botrytis cinerea</i> )	200 mL/100 L water	Spray at 14 day intervals commencing when the disease first becomes apparent and continuing until conditions no longer favour the disease. Spraying Saintpaulia, Poinsettia and Cyclamen flowers may result in some petal scorch. Tepid water should be used and wet plants protected from direct sunlight. This use is subject to an AFRAC anti-resistance strategy: Do not apply more than 3 consecutive Chipco Rovral GT or related dicarboximide fungicides.
Recreational Turf	Brown Patch ( <i>Rhizoctonia solani</i> )	120 to 180 mL /10 to 150 L water /100 m <sup>2</sup>	<u>Curative Program:</u> Repeat treatment after 14 days. Use the higher rate under conditions of severe disease pressure. <u>For preventative treatment:</u> Apply as a monthly spray program. This use is subject to an AFRAC anti-resistance strategy. Do not apply more than 2 consecutive sprays of Chipco Rovral GT or related dicarboximide fungicides unless tank mixed with a fungicide from a different activity group.
	Dollar Spot ( <i>Sclerotinia homoeocarpa</i> )	120 mL/10 to 150 L of water/100 m <sup>2</sup>	<u>For Curative Treatment:</u> Repeat treatment in 14 days. <u>For Preventative Treatment:</u> Apply monthly. This use is subject to an AFRAC anti-resistance strategy. Do not apply more than two consecutive sprays of Chipco Rovral GT or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Fusarium Patch ( <i>Fusarium nivale</i> ) (NOT Qld)	180 mL/10 to 150 L of water/100 m <sup>2</sup>	<u>For Preventative Treatment only:</u> Apply as a monthly soil drench throughout the year. Water in thoroughly after application. This use is subject to an AFRAC anti-resistance strategy. Do not apply more than two consecutive sprays of Chipco Rovral GT or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Spring Dead Spot ( <i>Leptosphaeria</i> spp.)	90 mL/10 to 150 L of water/100 m <sup>2</sup>	<u>For Curative Treatment:</u> Apply as a foliar spray and repeat application after 7 to 14 days. For white spot symptoms on couch grass ( <i>Cynodon dactylon</i> ) only, use higher water volumes of 100 to 150L of water per 100m <sup>2</sup> . <u>For Preventative Treatment:</u> Apply as a high volume foliar spray. Repeat applications at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. Do not apply more than two consecutive sprays of Chipco Rovral GT or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Helminthosporium disease affecting leaf, crown and roots: (Black and white Helminthosporium spot <i>Bipolaris</i> spp. <i>Drechslera</i> spp. <i>Exserohilum</i> spp.)		

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**