

CARTON FRONT PANEL

READ SAFETY DIRECTIONS BEFORE OPENING OR USING**Maxforce[®] White****COCKROACH GEL****ACTIVE CONSTITUENT: 21.5 g/kg IMIDACLOPRID****GROUP****4A****INSECTICIDE****30 g
NET**

For the control of cockroaches in the situations specified in the directions for use.

The syringe must not be sold separately. Do not destroy carton while product still remains.(label code)

CARTON REAR PANEL**DIRECTIONS FOR USE**

Situations	Pest	Rate	Critical Comments
Barracks, Farm buildings, Food processing establishments, Factories, Homes, Hospitals, Meatworks (non-product areas), Offices, Restaurants, Schools, Shops, Storerooms, Transport vehicles (including aircrafts) & facilities.	<u>Small cockroaches</u> e.g. German Cockroach (<i>Blattella germanica</i>)	<u>Light infestations</u> 1 placement per m ² <u>Heavy infestations</u> 2 placements per m ² 1 placement = a 5 mm ball (about the size of a small pea or approx. 0.1 g) or a line about 2.5cm long	Apply gel in out of the way locations where cockroaches occur e.g., under fridges, near water heaters, behind microwave ovens, garbage bins etc. Re-assessment of treated areas one month after initial baiting is recommended to evaluate if additional baiting is required. Bait placements should be replenished when consumed or at least every three months as long as pest populations persist.
	<u>Large cockroaches</u> e.g. American Cockroach (<i>Periplaneta</i> spp.)	<u>Light infestations</u> 2 placements per m ² <u>Heavy infestations</u> 3 placements per m ²	<i>Periplaneta</i> spp. require more bait because of the quantity which can be consumed by a single cockroach. Reassess baiting effectiveness as above.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Do not allow to contact food, livestock feed or their water. Do not place gel on surfaces where food is prepared or on utensils that are used for food preparation, on food packing materials or on surfaces where food packing materials are stored.

GENERAL INSTRUCTIONS:

Apply using suitable applicator such as BAITGUN®. Gel should be applied out of reach of children, pets and other animals. Do not place gel on surfaces which are regularly washed.

Insecticide Resistance Warning

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

For insecticide resistance management, Maxforce White Cockroach Gel Residual Insecticide is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Maxforce White and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Maxforce White or other Group 4A insecticides are used repeatedly. The effectiveness of Maxforce White on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of Maxforce White to control resistant insects. Maxforce White may be subject to specific resistance management strategies. For further information contact your local supplier or Bayer Environmental Science representative.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Do NOT contaminate streams, rivers or waterways with the gel or used syringes.

CARTON REAR PANEL CONT.

STORAGE AND DISPOSAL

Keep out of reach of children. Do not store in direct sunlight. Store in the closed, original carton, in a cool dry place. Completely empty syringe in normal use. Dispose of empty syringe and carton by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Suitable for use in export registered meat establishments.

APVMA Approval Number: 61937/53403

Maxforce® is a Registered Trademark of Bayer.

Bayer CropScience Pty Ltd

A.B.N. 87 000 226 022

391-393 Tooronga Rd

East Hawthorn Vic 3123

Phone: 03 9248-6888

Website: www.bayeres.com.au

Technical enquiries 1800 804 479

For 24 hour specialist advice in an emergency only phone 1800 033 111

Batch Number:

Date of Manufacture:

Bar code

Label code

SYRINGE LABEL – COMMON TO AU AND NZ**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

			
Maxforce[®] White			
COCKROACH GEL			
ACTIVE CONSTITUENT: 21.5 g/kg IMIDACLOPRID			
GROUP	4A	INSECTICIDE	

For the control of cockroaches in the situations specified in the Directions for Use.

30 g NET

Before using product read directions on the outer pack. Sale of this syringe except in outer carton is illegal. Do not destroy carton while product still remains.

STORAGE AND DISPOSAL

Keep out of reach of children. Do not destroy carton while product still remains. Do not store in direct sunlight. Store in the closed, original carton, in a cool dry place. Completely empty syringe in normal use. Dispose of empty syringe and carton by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone **Australia** 13 11 26, **NZ** 0800 764 766).

APVMA Approval No. 61937/53403

Maxforce[®] is a Registered Trademark of Bayer

Registered to:

Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn Vic. 3123
Phone Australia: (03) 9248 6888
Freecall NZ: 0800 227 012
www.bayeres.com.au

In NZ

APPROVED PURSUANT TO THE HSNO ACT 1996, No. HSR000676

See www.epa.govt.nz for approval controls

HSNO CLASSIFICATIONS:
6.9B, 9.1B, 9.2C, 9.4A

